



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Botrychium Jenmani in Cuba

R. C. BENEDICT

The genus *Botrychium* is very widespread in the tropics as well as in temperate regions, but, to judge from herbarium collections, in the tropics, the species are uncommon. *B. Jenmani* Und., as originally described and as treated in the North American Flora (Vol. 16: part 1), was based entirely on Jamaican material. Now, however, Cuba may be added to its range by reason of a recent collection made by Dr. J. A. Shafer, in February of this year, between La Perla and Santa Ana, in the province of Oriente. (*Shafer no. 8626*, Feb. 11, 1911.)

The plants collected were just in a mature fruiting condition, some of the sporangia having opened, the others still closed but full-sized. In Jamaica the fruiting period appears to range from late December to April, just about six months earlier than the fruiting season of *B. Underwoodianum* Maxon, the related Jamaican form. These two species form one of the several interesting pairs of *Botrychium* species, which occur in widely separated regions, and include in each locality two forms more or less alike in form and cutting but differing in the time of fruiting.

In Jamaica the pair comprises, as just noted, *Botrychium Jenmani* and *B. Underwoodianum*. In the southern United States a similar pair exists in the case of *B. biternatum* (Lam.) Und., and *B. alabamense* Maxon, fruiting in the early spring (February to April) and in the early fall (September), respectively. In California, *B. silaifolium* Presl and *B. californicum* Und. appear to constitute a similar pair, with *B. californicum* the earlier species. Now recently it has been suggested by a writer in the Fern Bulletin that there is here in the Eastern States a similar difference between *B. silaifolium intermedium* and *B.*

obliquum, the latter being said to be at least a month later in the same locality.

The explanation of the relationships of the extremely puzzling forms in the *Botrychium ternatum* group, here recognized as species, is not a simple one from our present knowledge of these plants. It is comparatively easy to arrange most of them in an evolutionary scheme, recognizing geographic distribution and consequent climatic difference as two chief causes of variation. This explanation, however, clears up only a part of the difficulty, and what remains is the more perplexing. More facts are needed regarding the habits, fruiting periods, and form variations. From the present knowledge, it appears to the writer that most of the distinct forms are deserving of the rank of species. It may be that new evidence, especially if discovered by experiment, will make a considerable change in our conception of these plants. The most promising field would appear to be along the line of pedigree cultures, but unfortunately it still remains to be discovered how these ferns may be raised from spore to adult condition.

COLUMBIA UNIVERSITY.